

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 January 2003 (23.01.2003)

PCT

(10) International Publication Number
WO 03/006666 A2

(51) International Patent Classification⁷: **C12P 13/00**

(21) International Application Number: **PCT/EP02/06187**

(22) International Filing Date: **6 June 2002 (06.06.2002)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
101 33 667.5 11 July 2001 (11.07.2001) DE
60/305,144 16 July 2001 (16.07.2001) US

(71) Applicant (*for all designated States except US*): **DE-GUSSA AG** [DE/DE]; Bennigsenplatz 1, 40474 Düsseldorf (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **SIEBELT, Nicole** [DE/DE]; Am Sennebach 28, 33397 Rietberg (DE). **WIDAWKA, Petra** [DE/DE]; Laerstrasse 12, 33615 Bielefeld (DE). **FARWICK, Mike** [DE/DE]; Gustav-Adolf-Strasse 11, 33615 Bielefeld (DE).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,

CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declaration under Rule 4.17:

— *of inventorship (Rule 4.17(iv)) for US only*

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/006666 A2

(54) Title: **PROCESS FOR THE PREPARATION OF L-AMINO ACIDS USING STRAINS OF THE ENTEROBACTERIACEAE FAMILY**

(57) Abstract: The invention relates to a process for the preparation of L-amino acids, in particular L-threonine, in which the following steps are carried out: a) fermentation of microorganisms of the Enterobacteriaceae family which produce the desired L-amino acid and in which at least one or more of the genes of cysteine biosynthesis chosen from the group consisting of cysG, cysB, cysZ, cysK, cysM, cysA, cysW, cysU, cysP, cysD, cysN, cysC, cysJ, cysI, cysH, cysE and sbp, or nucleotide sequences which code for these, is (are) enhanced, in particular over-expressed, b) concentration of the desired L-amino acid in the medium or in the cells of the bacteria, and c) isolation of the desired L-amino acid.

**Process for the Preparation of L-Amino Acids Using
Strains of the Enterobacteriaceae Family**

Field of the Invention

This invention relates to a process for the preparation of
5 L-amino acids, in particular L-threonine, using strains of
the Enterobacteriaceae family in which at least one or more
of the genes of the cysteine biosynthesis pathway (cysteine
biosynthetic pathway) chosen from the group consisting of
cysG, cysB, cysZ, cysK, cysM, cysA, cysW, cysU, cysP, cysD,
10 cysN, cysC, cysJ, cysI, cysH, cysE and sbp is (are)
enhanced.

Prior Art

L-Amino acids, in particular L-threonine, are used in human
medicine and in the pharmaceuticals industry, in the
15 foodstuffs industry and very particularly in animal
nutrition.

It is known to prepare L-amino acids by fermentation of
strains of Enterobacteriaceae, in particular Escherichia
coli (E. coli) and Serratia marcescens. Because of their
20 great importance, work is constantly being undertaken to
improve the preparation processes. Improvements to the
process can relate to fermentation measures, such as e.g.
stirring and supply of oxygen, or the composition of the
nutrient media, such as e.g. the sugar concentration during
25 the fermentation, or the working up to the product form, by
e.g. ion exchange chromatography, or the intrinsic output
properties of the microorganism itself.

Methods of mutagenesis, selection and mutant selection are
used to improve the output properties of these
30 microorganisms. Strains which are resistant to
antimetabolites, such as e.g. the threonine analogue α -
amino- β -hydroxyvaleric acid (AHV), or are auxotrophic for
metabolites of regulatory importance and produce L-amino

acids, such as e.g. L-threonine, are obtained in this manner.

Methods of the recombinant DNA technique have also been employed for some years for improving the strain of
5 strains of the Enterobacteriaceae family which produce L-amino acids, by amplifying individual amino acid biosynthesis genes and investigating the effect on the production.

Object of the Invention

- 10 The object of the invention is to provide new measures for improved fermentative preparation of L-amino acids, in particular L-threonine.

Summary of the Invention

- The invention provides a process for the fermentative
15 preparation of L-amino acids, in particular L-threonine, using microorganisms of the Enterobacteriaceae family which in particular already produce L-amino acids and in which at least one or more of the nucleotide sequence(s) which code(s) for the genes of the cysteine biosynthesis pathway
20 (cysteine biosynthetic pathway) chosen from the group consisting of cysG, cysB, cysZ, cysK, cysM, cysA, cysW, cysU, cysP, cysD, cysN, cysC, cysJ, cysI, cysH, cysE and sbp is (are) enhanced.

- The process according to the invention for the preparation
25 of amino acids comprises the following steps:

- a) fermentation of the microorganisms of the Enterobacteriaceae family which produce the desired L-amino acid and in which one or more of the genes of the cysteine biosynthesis pathway chosen from
30 the group consisting of cysG, cysB, cysZ, cysK, cysM, cysA, cysW, cysU, cysP, cysD, cysN, cysC, cysJ, cysI, cysH, cysE and sbp, or nucleotide

sequences which code for them is/are enhanced, in particular over-expressed,

- b) concentration of the desired L-amino acid in the medium or in the cells of the microorganisms, and
- 5 c) isolation of the desired L-amino acid, constituents of the fermentation broth and/or the biomass in its entirety or portions (> 0 to 100%) thereof optionally remaining in the product.

Detailed Description of the Invention

- 10 The use of endogenous genes is preferred. "Endogenous genes" or "endogenous nucleotide sequences" are understood as meaning the genes or nucleotide sequences present in the population of a species.

Where L-amino acids or amino acids are mentioned in the following, this means one or more amino acids, including their salts, chosen from the group consisting of L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine. L-Threonine is particularly preferred.

The term "enhancement" in this connection describes the increase in the intracellular activity or concentration of one or more enzymes or proteins in a microorganism which are coded by the corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or a gene or allele which codes for a corresponding enzyme or protein with a high activity, and optionally combining these measures.

- 30 By enhancement measures, in particular over-expression, the activity or concentration of the corresponding protein is in general increased by at least 10%, 25%, 50%, 75%, 100%,

150%, 200%, 300%, 400% or 500%, up to a maximum of 1000% or 2000%, based on that of the wild-type protein or the activity or concentration of the protein in the starting microorganism.

5 The process comprises carrying out the following steps:

- 10 a) fermentation of microorganisms of the Enterobacteriaceae family in which one or more of the genes of the cysteine biosynthesis pathway chosen from the group consisting of cysG, cysB, cysZ, cysK, cysM, cysA, cysW, cysU, cysP, cysD, cysN, cysC, cysJ, cysI, cysH, cysE and sbp is (are) enhanced,
 - 15 b) concentration of the corresponding L-amino acid in the medium or in the cells of the microorganisms of the Enterobacteriaceae family, and
 - c) isolation of the desired L-amino acid, constituents of the fermentation broth and/or the biomass in its entirety or portions (> 0 to 100 %) thereof optionally remaining in the product.
- 20 The microorganisms which the present invention provides can produce L-amino acids from glucose, sucrose, lactose, fructose, maltose, molasses, optionally starch, optionally cellulose or from glycerol and ethanol. They are representatives of the Enterobacteriaceae family chosen
- 25 from the genera Escherichia, Erwinia, Providencia and Serratia. The genera Escherichia and Serratia are preferred. Of the genus Escherichia the species Escherichia coli and of the genus Serratia the species Serratia marcescens are to be mentioned in particular.
- 30 Suitable strains, which produce L-threonine in particular, of the genus Escherichia, in particular of the species Escherichia coli, are, for example

- Escherichia coli TF427
- Escherichia coli H4578
- Escherichia coli KY10935
- Escherichia coli VNIIGenetika MG442
- 5 Escherichia coli VNIIGenetika M1
- Escherichia coli VNIIGenetika 472T23
- Escherichia coli BKIIM B-3996
- Escherichia coli kat 13
- Escherichia coli KCCM-10132

- 10 Suitable L-threonine-producing strains of the genus Serratia, in particular of the species Serratia marcescens, are, for example

- Serratia marcescens HNr21
- Serratia marcescens TLr156
- 15 Serratia marcescens T2000

- Strains from the Enterobacteriaceae family which produce L-threonine preferably have, inter alia, one or more genetic or phenotypic features chosen from the group consisting of:
- 20 resistance to α -amino- β -hydroxyvaleric acid, resistance to thialysine, resistance to ethionine, resistance to α -methylserine, resistance to diaminosuccinic acid, resistance to α -aminobutyric acid, resistance to borrelidin, resistance to rifampicin, resistance to valine analogues, such as, for example, valine hydroxamate,
 - 25 resistance to purine analogues, such as, for example, 6-dimethylaminopurine, a need for L-methionine, optionally a partial and compensable need for L-isoleucine, a need for meso-diaminopimelic acid, auxotrophy in respect of threonine-containing dipeptides, resistance to L-threonine,
 - 30 resistance to L-homoserine, resistance to L-lysine, resistance to L-methionine, resistance to L-glutamic acid, resistance to L-aspartate, resistance to L-leucine, resistance to L-phenylalanine, resistance to L-serine, resistance to L-cysteine, resistance to L-valine,
 - 35 sensitivity to fluoropyruvate, defective threonine

- dehydrogenase, optionally an ability for sucrose utilization, enhancement of the threonine operon, enhancement of homoserine dehydrogenase I-aspartate kinase I, preferably of the feed back resistant form, enhancement
 5 of homoserine kinase, enhancement of threonine synthase, enhancement of aspartate kinase, optionally of the feed back resistant form, enhancement of aspartate semialdehyde dehydrogenase, enhancement of phosphoenol pyruvate carboxylase, optionally of the feed back resistant form,
 10 enhancement of phosphoenol pyruvate synthase, enhancement of transhydrogenase, enhancement of the RhtB gene product, enhancement of the RhtC gene product, enhancement of the YfiK gene product, enhancement of a pyruvate carboxylase, and attenuation of acetic acid formation.
- 15 It has been found that microorganisms of the Enterobacteriaceae family produce L-amino acids, in particular L-threonine, in an improved manner after enhancement, in particular over-expression, of at least one or more of the genes of the cysteine biosynthesis pathway
 20 chosen from the group consisting of cysG, cysB, cysZ, cysK, cysM, cysA, cysW, cysU, cysP, cysD, cysN, cysC, cysJ, cysI, cysH, cysE and sbp.

The nucleotide sequences of the genes of *Escherichia coli* belong to the prior art (See following text references) and
 25 can also be found in the genome sequence of *Escherichia coli* published by Blattner et al. (Science 277: 1453 - 1462 (1997)). The genes and activities of the cysteine biosynthesis pathway (cysteine biosynthetic pathway) are also described in summary form in Kredich (In: Neidhardt
 30 (ed), *Escherichia coli* and *Salmonella*, American Society for Microbiology, Washington, D.C., USA: 514-527 (1996)).

cysG gene:

Description: Uroporphyrinogen III C methyl-transferase;
 precorrin-2 oxidase; ferrochelatase

35 EC No.: 2.1.1.107; 1.-.-.-; 4.99.1.-

Reference: Peakman et al.; European Journal of
Biochemistry 191(2): 315-323 (1990)
Macdonald and Cole; Molecular and General
Genetics 200(2): 328-334 (1985) Warren et
5 al.; Biochemical Journal 265(3):725-729
(1990) Spencer et al.; FEBS Letters 335(1):
57-60 (1993)

Accession No.: AE000412

cysB gene:

10 Description: Positive regulator of the cys regulon,
transcription activator

Reference: Ostrowski et al.; Journal of Biological
Chemistry 262(13): 5999-6005 (1987)
Mascarenhas and Yudkin; Molecular and
15 General Genetics 177(3): 535-539 (1980)
Lochowska et al.; Journal of Biological
Chemistry 276(3): 2098-2107 (2001)

Accession No.: AE000225

cysZ gene:

20 Description: Sulfate transporter

Reference: Byrne et al.; Journal of Bacteriology
170(7): 3150-3157 (1988)

Accession No.: AE000329

cysK gene:

25 Description: Cysteine synthase A, O-acetylserine
(thiol)-lyase A

EC No.: 4.2.99.8

Reference: Byrne et al.; Journal of Bacteriology
170(7): 3150-3157 (1988) Boronat et al.;
30 Journal of General Microbiology 130: 673-
685 (1984) Levy and Danchin; Molecular
Microbiology 2(6): 777-783 (1988)

Accession No.: AE000329

Alternative gene name: cysZ

- cysM gene:
Description: Cysteine synthase B, O-acetylserine
(thiol)-lyase B
EC No.: 4.2.99.8
5 Reference: Sirko et al.; Journal of Bacteriology
172(6): 3351-3357 (1990) Sirko et al.;
Journal of General Microbiology 133: 2719-
2725 (1987)
Accession No.: AE000329
- 10 cysA gene:
Description: ATP-binding protein of the sulfate
transport system
Reference: Sirko et al.; Journal of Bacteriology
172(6): 3351-3357 (1990) Sirko et al.;
15 Journal of General Microbiology 133: 2719-
2725 (1987)
Accession No.: AE000329
- cysW gene:
Description: Membrane-bound sulfate transport protein
20 Reference: Sirko et al.; Journal of Bacteriology
172(6): 3351-3357 (1990)
Accession No.: AE000329, AE000330
- cysU gene:
Description: Permease protein of the sulfate transport
25 system
Reference: Sirko et al.; Journal of Bacteriology
172(6): 3351-3357 (1990) Hryniewicz et al.;
Journal of Bacteriology 172(6): 3358-3366
(1990)
30 Accession No.: AE000330
Alternative gene name: cyst
- cysP gene:
Description: Periplasmic thiosulfate-binding protein

- Reference: Hryniewicz et al.; Journal of Bacteriology
172(6): 3358-3366 (1990) Sirko et al.;
Journal of Bacteriology 177(14): 4134-4136
(1995)
- 5 Accession No.: AE000330
- cysD gene:
- Description: Sub-unit 2 of ATP sulfurylase (ATP:sulfate
adenylyl-transferase)
- EC No.: 2.7.7.4
- 10 Reference: Leyh et al.; Journal of Biological
Chemistry 267(15): 10405-10410 (1992) Leyh
et al.; Journal of Biological Chemistry
263(5): 2409-2416 (1988)
- Accession No.: AE000358
- 15 cysN gene:
- Description: Sub-unit 1 of ATP sulfurylase (ATP:sulfate
adenylyl-transferase)
- EC No.: 2.7.7.4
- Reference: Leyh et al.; Journal of Biological
20 Chemistry 267(15): 10405-10410 (1992) Leyh
et al.; Journal of Biological Chemistry
263(5): 2409-2416 (1988) Leyh and Suo;
Journal of Biological Chemistry 267(1):
542-545 (1992)
- 25 Accession No.: AE000358
- cysC gene:
- Description: Adenylyl sulfate kinase (APS kinase)
- EC No.: 2.7.1.25
- Reference: Leyh et al.; Journal of Biological
30 Chemistry 267(15): 10405-10410 (1992) Leyh
et al.; Journal of Biological Chemistry
263(5): 2409-2416 (1988)
- Accession No.: AE000358

cysJ gene:

Description: Flavoprotein of NADPH sulfite reductase

EC No.: 1.8.1.2

Reference: Ostrowski et al.; Journal of Biological
5 Chemistry 264(27): 15796-15808 (1989) Li et
al.; Gene 53(2-3): 227-234 (1987) Gaudu and
Fontecave; European Journal of Biochemistry
226(2): 459-463 (1994) Eschenbrenner et
10 al.; Journal of Biological Chemistry
270(35): 20550-20555 (1995)

Accession No.: AE000360

Alternative gene name: cysP

cysI gene:

Description: Haemoprotein of NADPH sulfite reductase

15 EC No.: 1.8.1.2

Reference: Ostrowski et al.; Journal of Biological
Chemistry 264(26): 15726-15737 (1989) Li et
al.; Gene 53(2-3): 227-234 (1987) Gaudu and
Fontecave; European Journal of Biochemistry
20 226(2): 459-463 (1994)

Accession No.: AE000360

Alternative gene name: cysQ

cysH gene:

Description: Phosphoadenosine phosphosulfate reductase
25 (PAPS reductase)

EC No.: 1.8.99.4

Reference: Ostrowski et al.; Journal of Biological
Chemistry 264(26): 15726-15737 (1989) Krone
et al.; Molecular and General Genetics
30 225(2): 314-319 (1991) Li et al.; Gene
53(2-3): 227-234 (1987) Berendt et al.;
European Journal of Biochemistry 233(1):
347-356 (1995)

Accession No.: AE000360

cysE gene:

Description: Serine acetyl-transferase

EC No.: 2.3.1.30

Reference: Denk and Böck; Journal of General

5 Microbiology 133, 515-25 (1987)

Accession No.: AE000438

sbp gene:

Description: Periplasmic sulfate-binding protein

Reference: Hellinga and Evans, European Journal of

10 Biochemistry 149(2): 363-373 (1985) Sirko
et al.; Journal of Bacteriology 177(14):
4134-4136 (1995) Jacobson et al.; Journal
of Biological Chemistry 266(8): 5220-5225
(1991)

15 Accession No.: AE000466

The nucleic acid sequences can be found in the databanks of
the National Center for Biotechnology Information (NCBI) of
the National Library of Medicine (Bethesda, MD, USA), the
nucleotide sequence databank of the European Molecular
20 Biologies Laboratories (EMBL, Heidelberg, Germany or
Cambridge, UK) or the DNA databank of Japan (DDBJ, Mishima,
Japan).

The genes described in the text references mentioned can be
used according to the invention. Alleles of the genes which
25 result from the degeneracy of the genetic code or due to
"sense mutations" of neutral function can furthermore be
used.

To achieve an enhancement, for example, expression of the
genes or the catalytic properties of the proteins can be
30 increased. The two measures can optionally be combined.

To achieve an over-expression, the number of copies of the
corresponding genes can be increased, or the promoter and
regulation region or the ribosome binding site upstream of

the structural gene can be mutated. Expression cassettes which are incorporated upstream of the structural gene act in the same way. By inducible promoters, it is additionally possible to increase the expression in the
5 course of fermentative L-threonine production. The expression is likewise improved by measures to prolong the life of the m-RNA. Furthermore, the enzyme activity is also enhanced by preventing the degradation of the enzyme protein. The genes or gene constructs can either be
10 present in plasmids with a varying number of copies, or can be integrated and amplified in the chromosome. Alternatively, an over-expression of the genes in question can furthermore be achieved by changing the composition of the media and the culture procedure.

15 Instructions in this context can be found by the expert, inter alia, in Chang and Cohen (Journal of Bacteriology 134: 1141-1156 (1978)), in Hartley and Gregori (Gene 13: 347-353 (1981)), in Amann and Brosius (Gene 40: 183-190 (1985)), in de Broer et al. (Proceedings of the National
20 Academy of Sciences of the United States of America 80: 21-25 (1983)), in LaVallie et al. (BIO/TECHNOLOGY 11: 187-193 (1993)), in PCT/US97/13359, in Llosa et al. (Plasmid 26: 222-224 (1991)), in Quandt and Klipp (Gene 80: 161-169 (1989)), in Hamilton (Journal of Bacteriology 171: 4617-
25 4622 (1989)), in Jensen and Hammer (Biotechnology and Bioengineering 58: 191-195 (1998)) and in known textbooks of genetics and molecular biology.

Plasmid vectors which are capable of replication in Enterobacteriaceae, such as e.g. cloning vectors derived
30 from pACYC184 (Bartolomé et al.; Gene 102: 75-78 (1991)), pTrc99A (Amann et al.; (Gene 69: 301-315 (1988)) or pSC101 derivatives (Vocke and Bastia, Proceedings of the National Academy of Sciences USA 80 (21): 6557-6561 (1983)) can be used. A strain transformed with a plasmid vector, where the
35 plasmid vector carries at least one or more of the genes

chosen from the group consisting of cysG, cysB, cysZ, cysK, cysM, cysA, cysW, cysU, cysP, cysD, cysN, cysC, cysJ, cysI, cysH, cysE and sbp or nucleotide sequences which code for these, can be employed in a process according to the
5 invention.

It is also possible to transfer mutations which affect the expression of the particular gene into various strains by sequence exchange (Hamilton et al. (Journal of Bacteriology 171: 4617 - 4622 (1989)), conjugation or transduction.

- 10 It may furthermore be advantageous for the production of L-amino acids, in particular L-threonine, with strains of the Enterobacteriaceae family to enhance one or more enzymes of the known threonine biosynthesis pathway or enzymes of anaplerotic metabolism or enzymes for the production of
15 reduced nicotinamide adenine dinucleotide phosphate or enzymes of glycolysis or PTS enzymes, in addition to enhancement of one or more of the genes of the cysteine biosynthesis pathway chosen from the group consisting of cysG, cysB, cysZ, cysK, cysM, cysA, cysW, cysU, cysP, cysD,
20 cysN, cysC, cysJ, cysI, cysH, cysE and sbp.

Thus, for example, at the same time one or more of the genes chosen from the group consisting of

- the thrABC operon which codes for aspartate kinase, homoserine dehydrogenase, homoserine kinase and
25 threonine synthase (US-A-4,278,765),
- the pyc gene of Corynebacterium glutamicum which codes for pyruvate carboxylase (WO 99/18228),
- the pps gene which codes for phosphoenol pyruvate synthase (Molecular and General Genetics 231: 332-336
30 (1992)),
- the ppc gene which codes for phosphoenol pyruvate carboxylase (Gene 31: 279-283 (1984)),

- the pntA and pntB genes which code for transhydrogenase (European Journal of Biochemistry 158: 647-653 (1986)),
- the rhtB gene which imparts homoserine resistance (EP-A-0 994 190),
- 5 • the mgo gene which codes for malate:quinone oxidoreductase (WO 02/06459),
- the rhtC gene which imparts threonine resistance (EP-A-1 013 765),
- the thrE gene of Corynebacterium glutamicum which codes for threonine export (WO 01/92545),
- 10 • the gdhA gene which codes for glutamate dehydrogenase (Nucleic Acids Research 11: 5257-5266 (1983); Gene 23: 199-209 (1983))
- the hns gene which codes for the DNA-binding protein HLP-II (Molecular and General Genetics 212: 199-202 (1988)),
- 15 • the pgm gene which codes for phosphoglucomutase (Journal of Bacteriology 176: 5847-5851 (1994)),
- the fba gene which codes for fructose biphosphate aldolase (Biochemical Journal 257: 529-534 (1989)),
- 20 • the ptsH gene of the ptsHIcrr operon which codes for the phosphohistidine protein hexose phosphotransferase of the phosphotransferase system PTS (Journal of Biological Chemistry 262: 16241-16253 (1987)),
- 25 • the ptsI gene of the ptsHIcrr operon which codes for enzyme I of the phosphotransferase system PTS (Journal of Biological Chemistry 262: 16241-16253 (1987)),
- the crr gene of the ptsHIcrr operon which codes for the glucose-specific IIA component of the phosphotransferase

system PTS (Journal of Biological Chemistry 262: 16241-16253 (1987)),

- the ptsG gene which codes for the glucose-specific IIBC component (Journal of Biological Chemistry 261: 16398-16403 (1986)),
 - the lrp gene which codes for the regulator of the leucine regulon (Journal of Biological Chemistry 266: 10768-10774 (1991)),
 - the mopB gene which codes for 10 Kd chaperone (Journal of Biological Chemistry 261: 12414-12419 (1986)) and is also known by the name groES,
 - the ahpC gene of the ahpCF operon which codes for the small sub-unit of alkyl hydroperoxide reductase (Proceedings of the National Academy of Sciences USA 92: 7617-7621 (1995))
 - the ahpF gene of the ahpCF operon which codes for the large sub-unit of alkyl hydroperoxide reductase (Proceedings of the National Academy of Sciences USA 92: 7617-7621 (1995))
- can be enhanced, in particular over-expressed.

It may furthermore be advantageous for the production of L-amino acids, in particular L-threonine, in addition to enhancement of one or more of the genes chosen from the group consisting of cysG, cysB, cysZ, cysK, cysM, cysA, cysW, cysU, cysP, cysD, cysN, cysC, cysJ, cysI, cysH, cysE and sbp, for one or more of the genes chosen from the group consisting of

- the tdh gene which codes for threonine dehydrogenase (Ravnikar and Somerville (Journal of Bacteriology 169: 4716-4721 (1987))),

- the mdh gene which codes for malate dehydrogenase (E.C. 1.1.1.37) (Vogel et al. (Archives in Microbiology 149: 36-42 (1987))),
- the gene product of the open reading frame (orf) yjfa
5 (Accession Number AAC77180 of the National Center for Biotechnology Information (NCBI, Bethesda, MD, USA)),
- the gene product of the open reading frame (orf) ytfP (Accession Number AAC77179 of the National Center for Biotechnology Information (NCBI, Bethesda, MD, USA)),
- 10 • the pckA gene which codes for the enzyme phosphoenolpyruvate carboxykinase (Medina et al. (Journal of Bacteriology 172: 7151-7156 (1990))),
- the poxB gene which codes for pyruvate oxidase (Grabau and Cronan (Nucleic Acids Research 14 (13), 5449-5460
15 (1986))),
- the aceA gene which codes for the enzyme isocitrate lyase (Matsuoko and McFadden (Journal of Bacteriology 170, 4528-4536 (1988))),
- the dgsA gene which codes for the DgsA regulator of the
20 phosphotransferase system (Hosono et al. (Bioscience, Biotechnology and Biochemistry 59: 256-251 (1995))) and is also known under the name of the mlc gene,
- the fruR gene which codes for the fructose repressor (Jahreis et al. (Molecular and General Genetics 226:
25 332-336 (1991))) and is also known by the name of the cra gene, and
- the rpoS gene which codes for the sigma³⁸ factor (WO 01/05939) and is also known under the name of the katF gene,
- 30 to be attenuated, in particular eliminated or for the expression thereof to be reduced.

The term "attenuation" in this connection describes the reduction or elimination of the intracellular activity or concentration of one or more enzymes or proteins in a microorganism which are coded by the corresponding DNA, for example by using a weak promoter or a gene or allele which codes for a corresponding enzyme or protein with a low activity or inactivates the corresponding enzyme or protein or gene, and optionally combining these measures.

By attenuation measures, the activity or concentration of the corresponding protein is in general reduced to 0 to 75%, 0 to 50%, 0 to 25%, 0 to 10% or 0 to 5% of the activity or concentration of the wild-type protein or of the activity or concentration of the protein in the starting microorganism.

It may furthermore be advantageous for the production of L-amino acids, in particular L-threonine, in addition to enhancement of one or more of the genes of the cysteine biosynthesis pathway chosen from the group consisting of cysG, cysB, cysZ, cysK, cysM, cysA, cysW, cysU, cysP, cysD, cysN, cysC, cysJ, cysI, cysH, cysE and sbp, to eliminate undesirable side reactions (Nakayama: "Breeding of Amino Acid Producing Microorganisms", in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

The microorganisms produced according to the invention can be cultured in the batch process (batch culture), the fed batch (feed process) or the repeated fed batch process (repetitive feed process). A summary of known culture methods is described in the textbook by Chmiel (Bioprozesstechnik 1. Einführung in die Bioverfahrenstechnik [Bioprocess Technology 1. Introduction to Bioprocess Technology (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen [Bioreactors and Peripheral Equipment] (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained in the handbook "Manual of Methods for General
5 Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981).

Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and optionally cellulose, oils and fats, such as e.g. soya oil, sunflower
10 oil, groundnut oil and coconut fat, fatty acids, such as e.g. palmitic acid, stearic acid and linoleic acid, alcohols, such as e.g. glycerol and ethanol, and organic acids, such as e.g. acetic acid, can be used as the source of carbon. These substances can be used individually or as
15 a mixture.

Organic nitrogen-containing compounds, such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium
20 phosphate, ammonium carbonate and ammonium nitrate, can be used as the source of nitrogen. The sources of nitrogen can be used individually or as a mixture.

Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-
25 containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of metals, such as e.g. magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be
30 employed in addition to the above-mentioned substances. Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH of the culture.

- 5 Antifoams, such as e.g. fatty acid polyglycol esters, can be employed to control the development of foam. Suitable substances having a selective action, e.g. antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain aerobic conditions, oxygen or
10 oxygen-containing gas mixtures, such as e.g. air, are introduced into the culture. The temperature of the culture is usually 25°C to 45°C, and preferably 30°C to 40°C. Culturing is continued until a maximum of L-amino acids or L-threonine has formed. This target is usually
15 reached within 10 hours to 160 hours.

The analysis of L-amino acids can be carried out by anion exchange chromatography with subsequent ninhydrin derivation, as described by Spackman et al. (Analytical Chemistry 30: 1190-1206 (1958)), or it can take place by
20 reversed phase HPLC as described by Lindroth et al. (Analytical Chemistry 51: 1167-1174 (1979)).

The process according to the invention is used for the fermentative preparation of L-amino acids, such as, for example, L-threonine, L-isoleucine, L-valine, L-methionine,
25 L-homoserine and L-lysine, in particular L-threonine.

The minimal (M9) and complete media (LB) for *Escherichia coli* used are described by J.H. Miller (A short course in bacterial genetics (1992), Cold Spring Harbor Laboratory Press). The isolation of plasmid DNA from *Escherichia coli*
30 and all techniques of restriction, ligation, Klenow and alkaline phosphatase treatment are carried out by the method of Sambrook et al. (Molecular cloning - A Laboratory Manual (1989) Cold Spring Harbor Laboratory Press). Unless described otherwise, the transformation of *Escherichia coli*
35 is carried out by the method of Chung et al. (Proceedings

of the National Academy of Sciences of the United States of America (1989) 86: 2172-2175).

The incubation temperature for the preparation of strains and transformants is 37°C.

5 Example 1

Preparation of L-threonine using the *cysB* gene

1a) Construction of the expression plasmid pTrc99AcysB

The *cysB* gene from *E. coli* K12 is amplified using the polymerase chain reaction (PCR) and synthetic
10 oligonucleotides. Starting from the nucleotide sequence of the *cysB* gene in *E. coli* K12 MG1655 (Accession Number AE000225, Blattner et al. (Science 277: 1453-1462 (1997))), PCR primers are synthesized (MWG Biotech, Ebersberg, Germany):

15 *cysB*1: 5' - GCGTCTAAGTGGATGGTTTAAC - 3' (SEQ ID No. 1)

*cysB*2: 5' - GGTGCCGAAAATAACGCAAG - 3' (SEQ ID No. 2)

The chromosomal *E. coli* K12 MG1655 DNA employed for the PCR is isolated according to the manufacturer's instructions with "Qiagen Genomic-tips 100/G" (QIAGEN, Hilden, Germany).
20 A DNA fragment approx. 1000 bp in size can be amplified with the specific primers under standard PCR conditions (Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press) with Pfu-DNA polymerase (Promega Corporation, Madison, USA). The PCR product is
25 ligated according to the manufacturer's instructions with the vector pCR-Blunt II-TOPO (Zero Blunt TOPO PCR Cloning Kit, Invitrogen, Groningen, The Netherlands) and transformed into the *E. coli* strain TOP10.

Selection of plasmid-carrying cells takes place on LB agar,
30 to which 50 µg/ml kanamycin are added. After isolation of the plasmid DNA, the vector pCR-Blunt II-TOPO-*cysB* is

cleaved with the restriction enzymes HindIII and XbaI and, after separation in 0.8% agarose gel, the cysB fragment is isolated with the aid of the QIAquick Gel Extraction Kit (QIAGEN, Hilden, Germany). The vector pTrc99A (Pharmacia Biotech, Uppsala, Sweden) is cleaved with the enzymes HindIII and XbaI and ligation is carried out with the cysB fragment isolated. The E. coli strain XL1-Blue MRF' (Stratagene, La Jolla, USA) is transformed with the ligation batch and plasmid-carrying cells are selected on LB agar, to which 50 µg/ml ampicillin are added. Successful cloning can be demonstrated after plasmid DNA isolation by control cleavage with the enzymes ScaI and SmaI. The plasmid is called pTrc99AcysB (Figure 1).

1b) Preparation of L-threonine with the strain
MG442/pTrc99AcysB

The L-threonine-producing E. coli strain MG442 is described in the patent specification US-A- 4,278,765 and deposited as CMIM B-1628 at the Russian National Collection for Industrial Microorganisms (VKPM, Moscow, Russia).

The strain MG442 is transformed with the expression plasmid pTrc99AcysB described in example Ia and with the vector pTrc99A and plasmid-carrying cells are selected on LB agar with 50 µg/ml ampicillin. The strains MG442/pTrc99AcysB and MG442/pTrc99A are formed in this manner. Selected individual colonies are then multiplied further on minimal medium with the following composition: 3.5 g/l Na₂HPO₄*2H₂O, 1.5 g/l KH₂PO₄, 1 g/l NH₄Cl, 0.1 g/l MgSO₄*7H₂O, 2 g/l glucose, 20 g/l agar, 50 mg/l ampicillin. The formation of L-threonine is checked in batch cultures of 10 ml contained in 100 ml conical flasks. For this, 10 ml of preculture medium of the following composition: 2 g/l yeast extract, 10 g/l (NH₄)₂SO₄, 1 g/l KH₂PO₄, 0.5 g/l MgSO₄*7H₂O, 15 g/l CaCO₃, 20 g/l glucose, 50 mg/l ampicillin are inoculated and the batch is incubated for 16 hours at 37°C and 180 rpm

on an ESR incubator from Kühner AG (Birsfelden, Switzerland).

250 µl portions of this preculture are transinoculated into 10 ml of production medium (25 g/l $(\text{NH}_4)_2\text{SO}_4$, 2 g/l KH_2PO_4 , 5 1 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.03 g/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.018 g/l $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 30 g/l CaCO_3 , 20 g/l glucose, 50 mg/l ampicillin) and the batch is incubated for 48 hours at 37°C. For complete induction of the expression of the *cysB* gene, 100 mg/l isopropyl β-D-thiogalactopyranoside (IPTG) are added in 10 parallel batches. The formation of L-threonine by the starting strain MG442 is investigated in the same manner, but no addition of ampicillin to the medium takes place. After the incubation the optical density (OD) of the culture suspension is determined with an LP2W photometer 15 from Dr. Lange (Düsseldorf, Germany) at a measurement wavelength of 660 nm.

The concentration of L-threonine formed is then determined in the sterile-filtered culture supernatant with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) 20 by ion exchange chromatography and post-column reaction with ninhydrin detection.

The result of the experiment is shown in Table 1.

Table 1

Strain	Additives	OD (660 nm)	L-Threonine g/l
MG442	-	5.6	1.4
MG442/pTrc99A	-	3.8	1.3
MG442/pTrc99AcysB	-	4.4	1.7
MG442/pTrc99AcysB	IPTG	5.4	2.0

Example 2

Preparation of L-threonine using the cysK gene

2a) Construction of the expression plasmid pTrc99AcysK

The cysK gene from E. coli K12 is amplified using the
5 polymerase chain reaction (PCR) and synthetic
oligonucleotides. Starting from the nucleotide sequence of
the cysK gene in E. coli K12 MG1655 (Accession Number
AE000329, Blattner et al. (Science 277: 1453-1462 (1997))),
PCR primers are synthesized (MWG Biotech, Ebersberg;
10 Germany):

cysK1: 5' - CAGTTAAGGACAGGCCATGAG - 3' (SEQ ID No. 3)

cysK2: 5' - GCTGGCATTACTGTTGCAATTC - 3' (SEQ ID No. 4)

The chromosomal E. coli K12 MG1655 DNA employed for the PCR
is isolated according to the manufacturer's instructions
15 with "Qiagen Genomic-tips 100/G" (QIAGEN, Hilden, Germany).
A DNA fragment approx. 1000 bp in size can be amplified
with the specific primers under standard PCR conditions
(Innis et al. (1990) PCR Protocols. A Guide to Methods and
Applications, Academic Press) with Pfu-DNA polymerase
20 (Promega Corporation, Madison, USA). The PCR product is
ligated according to the manufacturer's instructions with
the vector pCR-Blunt II-TOPO (Zero Blunt TOPO PCR Cloning
Kit, Invitrogen, Groningen, The Netherlands) and
transformed into the E. coli strain TOP10.

25 Selection of plasmid-carrying cells takes place on LB agar,
to which 50 µg/ml kanamycin are added. After isolation of
the plasmid DNA, the vector pCR-Blunt II-TOPO-cysK is
cleaved with the restriction enzymes SpeI and XbaI and,
after separation in 0.8% agarose gel, the cysK fragment is
30 isolated with the aid of the QIAquick Gel Extraction Kit
(QIAGEN, Hilden, Germany). The vector pTrc99A (Pharmacia
Biotech, Uppsala, Sweden) is cleaved with the enzyme XbaI

and ligation is carried out with the cysK fragment isolated. The E. coli strain XL1-Blue MRF' (Stratagene, La Jolla, USA) is transformed with the ligation batch and plasmid-carrying cells are selected on LB agar, to which
5 50 µg/ml ampicillin are added. Successful cloning can be demonstrated after plasmid DNA isolation by control cleavage with the enzymes HindIII and PvuII. The plasmid is called pTrc99AcysK (Figure 2).

2b) Preparation of L-threonine with the strain
10 MG442/pTrc99AcysK

The L-threonine-producing E. coli strain MG442 is described in the patent specification US-A- 4,278,765 and deposited as CMIM B-1628 at the Russian National Collection for Industrial Microorganisms (VKPM, Moscow, Russia).

15 The strain MG442 is transformed with the expression plasmid pTrc99AcysK described in example 2a and with the vector pTrc99A and plasmid-carrying cells are selected on LB agar with 50 µg/ml ampicillin. The strains MG442/pTrc99AcysK and MG442/pTrc99A are formed in this manner. Selected
20 individual colonies are then multiplied further on minimal medium with the following composition: 3.5 g/l $\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$, 1.5 g/l KH_2PO_4 , 1 g/l NH_4Cl , 0.1 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 2 g/l glucose, 20 g/l agar, 50 mg/l ampicillin. The formation of L-threonine is checked in batch cultures of 10 ml contained
25 in 100 ml conical flasks. For this, 10 ml of preculture medium of the following composition: 2 g/l yeast extract, 10 g/l $(\text{NH}_4)_2\text{SO}_4$, 1 g/l KH_2PO_4 , 0.5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 15 g/l CaCO_3 , 20 g/l glucose, 50 mg/l ampicillin are inoculated and the batch is incubated for 16 hours at 37°C and 180 rpm
30 on an ESR incubator from Kühner AG (Birsfelden, Switzerland).

250 µl portions of this preculture are transinoculated into 10 ml of production medium (25 g/l $(\text{NH}_4)_2\text{SO}_4$, 2 g/l KH_2PO_4 , 1 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.03 g/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.018 g/l $\text{MnSO}_4 \cdot 1\text{H}_2\text{O}$,

30 g/l CaCO₃, 20 g/l glucose, 50 mg/l ampicillin) and the batch is incubated for 48 hours at 37°C. The formation of L-threonine by the starting strain MG442 is investigated in the same manner, but no addition of ampicillin to the medium takes place. After the incubation the optical density (OD) of the culture suspension is determined with an LP2W photometer from Dr. Lange (Düsseldorf, Germany) at a measurement wavelength of 660 nm.

The concentration of L-threonine formed is then determined in the sterile-filtered culture supernatant with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column reaction with ninhydrin detection.

The result of the experiment is shown in Table 2.

15

Table 2

Strain	OD (660 nm)	L-Threonine g/l
MG442	5.6	1.4
MG442/pTrc99A	3.8	1.3
MG442/pTrc99AcysK	5.6	2.1

Example 3

Preparation of L-threonine using the cysM gene

3a) Construction of the expression plasmid pTrc99AcysM

20 The cysM gene from E. coli K12 is amplified using the polymerase chain reaction (PCR) and synthetic oligonucleotides. Starting from the nucleotide sequence of the cysM gene in E. coli K12 MG1655 (Accession Number

AE000329, Blattner et al. (Science 277: 1453-1462 (1997)), PCR primers are synthesized (MWG Biotech, Ebersberg, Germany). The sequences of the primers are modified such that recognition sites for restriction enzymes are formed.

- 5 The recognition sequence for XbaI is chosen for the cysM1 primer and the recognition sequence for HindIII for the cysM2 primer, which are marked by underlining in the nucleotide sequence shown below:

cysM1: 5' - CGCATCAGTCTAGACCACGTTAGGATAG - 3'
10 (SEQ ID No. 5)

cysM2: 5' - CATCAGTCTCCGAAGCTTTTAATCC - 3'
(SEQ ID No. 6)

- The chromosomal E. coli K12 MG1655 DNA employed for the PCR is isolated according to the manufacturer's instructions
15 with "Qiagen Genomic-tips 100/G" (QIAGEN, Hilden, Germany). A DNA fragment approx. 950 bp in size can be amplified with the specific primers under standard PCR conditions (Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press) with Pfu-DNA polymerase
20 (Promega Corporation, Madison, USA). The PCR product is ligated according to the manufacturer's instructions with the vector pCR-Blunt II-TOPO (Zero Blunt TOPO PCR Cloning Kit, Invitrogen, Groningen, The Netherlands) and transformed into the E. coli strain TOP10.

- 25 Selection of plasmid-carrying cells takes place on LB agar, to which 50 µg/ml kanamycin are added. After isolation of the plasmid DNA, the vector pCR-Blunt II-TOPO-cysM is cleaved with the restriction enzymes HindIII and XbaI and, after separation in 0.8% agarose gel, the cysM fragment is
30 isolated with the aid of the QIAquick Gel Extraction Kit (QIAGEN, Hilden, Germany). The vector pTrc99A (Pharmacia Biotech, Uppsala, Sweden) is cleaved with the enzymes HindIII and XbaI and ligation is carried out with the cysM fragment isolated. The E. coli strain XL1-Blue MRF'

(Stratagene, La Jolla, USA) is transformed with the ligation batch and plasmid-carrying cells are selected on LB agar, to which 50 µg/ml ampicillin are added. Successful cloning can be demonstrated after plasmid DNA isolation by control cleavage with the enzymes EcoRV, Eco91I, PstI and SspI. The plasmid is called pTrc99AcysM (Figure 3).

3b) Preparation of L-threonine with the strain MG442/pTrc99AcysM

The L-threonine-producing E. coli strain MG442 is described in the patent specification US-A- 4,278,765 and deposited as CMIM B-1628 at the Russian National Collection for Industrial Microorganisms (VKPM, Moscow, Russia).

The strain MG442 is transformed with the expression plasmid pTrc99AcysM described in example 3a and with the vector pTrc99A and plasmid-carrying cells are selected on LB agar with 50 µg/ml ampicillin. The strains MG442/pTrc99AcysM and MG442/pTrc99A are formed in this manner. Selected individual colonies are then multiplied further on minimal medium with the following composition: 3.5 g/l $\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$, 1.5 g/l KH_2PO_4 , 1 g/l NH_4Cl , 0.1 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 2 g/l glucose, 20 g/l agar, 50 mg/l ampicillin. The formation of L-threonine is checked in batch cultures of 10 ml contained in 100 ml conical flasks. For this, 10 ml of preculture medium of the following composition: 2 g/l yeast extract, 10 g/l $(\text{NH}_4)_2\text{SO}_4$, 1 g/l KH_2PO_4 , 0.5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 15 g/l CaCO_3 , 20 g/l glucose, 50 mg/l ampicillin are inoculated and the batch is incubated for 16 hours at 37°C and 180 rpm on an ESR incubator from Kühner AG (Birsfelden, Switzerland).

250 µl portions of this preculture are transinoculated into 10 ml of production medium (25 g/l $(\text{NH}_4)_2\text{SO}_4$, 2 g/l KH_2PO_4 , 1 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.03 g/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.018 g/l $\text{MnSO}_4 \cdot 1\text{H}_2\text{O}$, 30 g/l CaCO_3 , 20 g/l glucose, 50 mg/l ampicillin) and the batch is incubated for 48 hours at 37°C. The formation of

L-threonine by the starting strain MG442 is investigated in the same manner, but no addition of ampicillin to the medium takes place. After the incubation the optical density (OD) of the culture suspension is determined with
5 an LP2W photometer from Dr. Lange (Düsseldorf, Germany) at a measurement wavelength of 660 nm.

The concentration of L-threonine formed is then determined in the sterile-filtered culture supernatant with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany)
10 by ion exchange chromatography and post-column reaction with ninhydrin detection.

The result of the experiment is shown in Table 3.

Table 3

Strain	OD (660 nm)	L-Threonine g/l
MG442	5.6	1.4
MG442/pTrc99A	3.8	1.3
MG442/pTrc99AcysM	1.6	2.0

15 Example 4

Preparation of L-threonine using the cysP, cysU, cysW and cysA genes

4a) Construction of the expression plasmid pTrc99AcysPUWA

The cysP, cysU, cysW and cysA genes from E. coli K12 are
20 amplified using the polymerase chain reaction (PCR) and synthetic oligonucleotides. Starting from the nucleotide sequence of the cysP, cysU, cysW and cysA genes in E. coli K12 MG1655 (Accession Number AE000329 and AE000330,

Blattner et al. (Science 277: 1453-1462 (1997)), PCR primers are synthesized (MWG Biotech, Ebersberg, Germany). The sequences of the primers are modified such that recognition sites for restriction enzymes are formed. The
5 recognition sequence for XbaI is chosen for the cysPUWA1 primer and the recognition sequence for HindIII for the cysPUWA2 primer, which are marked by underlining in the nucleotide sequence shown below:

cysPUWA1: 5' - GTCTCTAGATAAATAAGGGTGC^{GAAT}GGC - 3'
10 (SEQ ID No. 7)

cysPUWA2: 5' - CCGGGCGTTTAAAGCTTCACTCAACC - 3'
(SEQ ID No. 8)

The chromosomal E. coli K12 MG1655 DNA employed for the PCR is isolated according to the manufacturer's instructions
15 with "Qiagen Genomic-tips 100/G" (QIAGEN, Hilden, Germany). A DNA fragment approx. 3900 bp in size can be amplified with the specific primers under standard PCR conditions (Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press) with Pfu-DNA polymerase
20 (Promega Corporation, Madison, USA). The PCR product is cleaved with the restriction enzymes XbaI and HindIII and ligated with the vector pTrc99A (Pharmacia Biotech, Uppsala, Sweden), which has been digested with the enzymes XbaI and HindIII. The E. coli strain XL1-Blue MRF'
25 (Stratagene, La Jolla, USA) is transformed with the ligation batch and plasmid-carrying cells are selected on LB agar, to which 50 µg/ml ampicillin are added. Successful cloning can be demonstrated after plasmid DNA isolation by control cleavage with the enzymes BamHI, EcoRV, MluI, NdeI
30 and SspI. The plasmid is called pTrc99AcysPUWA (Figure 4).

4b) Preparation of L-threonine with the strain
MG442/pTrc99AcysPUWA

The L-threonine-producing *E. coli* strain MG442 is described in the patent specification US-A- 4,278,765 and deposited
5 as CMIM B-1628 at the Russian National Collection for Industrial Microorganisms (VKPM, Moscow, Russia).

The strain MG442 is transformed with the expression plasmid pTrc99AcysPUWA described in example 4a and with the vector pTrc99A and plasmid-carrying cells are selected on LB agar
10 with 50 µg/ml ampicillin. The strains MG442/pTrc99AcysPUWA and MG442/pTrc99A are formed in this manner. Selected individual colonies are then multiplied further on minimal medium with the following composition: 3.5 g/l Na₂HPO₄*2H₂O, 1.5 g/l KH₂PO₄, 1 g/l NH₄Cl, 0.1 g/l MgSO₄*7H₂O, 2 g/l
15 glucose, 20 g/l agar, 50 mg/l ampicillin. The formation of L-threonine is checked in batch cultures of 10 ml contained in 100 ml conical flasks. For this, 10 ml of preculture medium of the following composition: 2 g/l yeast extract, 10 g/l (NH₄)₂SO₄, 1 g/l KH₂PO₄, 0.5 g/l MgSO₄*7H₂O, 15 g/l
20 CaCO₃, 20 g/l glucose, 50 mg/l ampicillin are inoculated and the batch is incubated for 16 hours at 37°C and 180 rpm on an ESR incubator from Kühner AG (Birsfelden, Switzerland).

250 µl portions of this preculture are transinoculated into
25 10 ml of production medium (25 g/l (NH₄)₂SO₄, 2 g/l KH₂PO₄, 1 g/l MgSO₄*7H₂O, 0.03 g/l FeSO₄*7H₂O, 0.018 g/l MnSO₄*1H₂O, 30 g/l CaCO₃, 20 g/l glucose, 50 mg/l ampicillin) and the batch is incubated for 48 hours at 37°C. For complete induction of the expression of the cysPUWA genes, 100 mg/l
30 isopropyl β-D-thiogalactopyranoside (IPTG) are added in parallel batches. The formation of L-threonine by the starting strain MG442 is investigated in the same manner, but no addition of ampicillin to the medium takes place. After the incubation the optical density (OD) of the
35 culture suspension is determined with an LP2W photometer

from Dr. Lange (Düsseldorf, Germany) at a measurement wavelength of 660 nm.

The concentration of L-threonine formed is then determined in the sterile-filtered culture supernatant with an amino
5 acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column reaction with ninhydrin detection.

The result of the experiment is shown in Table 4.

Table 4

Strain	Additives	OD (660 nm)	L-Threonine g/l
MG442	-	5.6	1.4
MG442/pTrc99A	-	3.8	1.3
MG442/pTrc99AcysPUWA	-	5.5	1.7
MG442/pTrc99AcysPUWA	IPTG	6.5	2.1

10

Example 5

Preparation of L-threonine using the cysD, cysN and cysC genes

5a) Construction of the expression plasmid pTrc99AcysDNC

15 The cysD, cysN and cysC genes from E. coli K12 are amplified using the polymerase chain reaction (PCR) and synthetic oligonucleotides. Starting from the nucleotide sequence of the cysD, cysN and cysC genes in E. coli K12 MG1655 (Accession Number AE000358, Blattner et al. (Science
20 277: 1453-1462 (1997))), PCR primers are synthesized (MWG Biotech, Ebersberg, Germany). The sequences of the primers

are modified such that recognition sites for restriction enzymes are formed. The recognition sequence for XbaI is chosen for the cysDNC1 primer and the recognition sequence for HindIII for the cysDNC2 primer, which are marked by underlining in the nucleotide sequence shown below:

cysDNC1: 5' - GCAAGAAAATAGCGGTCTAGATAAGGAACG - 3'
(SEQ ID No. 9)

cysDNC2: 5' - CATGGAAAGCTTGTGGTGTCTCAGG - 3'
(SEQ ID No. 10)

- 10 The chromosomal E. coli K12 MG1655 DNA employed for the PCR is isolated according to the manufacturer's instructions with "Qiagen Genomic-tips 100/G" (QIAGEN, Hilden, Germany). A DNA fragment approx. 3000 bp in size can be amplified with the specific primers under standard PCR conditions
- 15 (Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press) with Pfu-DNA polymerase (Promega Corporation, Madison, USA). The PCR product is cleaved with the restriction enzymes XbaI and HindIII and ligated with the vector pTrc99A (Pharmacia Biotech,
- 20 Uppsala, Sweden), which has been digested with the enzymes XbaI and HindIII. The E. coli strain XL1-Blue MRF' (Stratagene, La Jolla, USA) is transformed with the ligation batch and plasmid-carrying cells are selected on LB agar, to which 50 µg/ml ampicillin are added. Successful
- 25 cloning can be demonstrated after plasmid DNA isolation by control cleavage with the enzymes EcoRV, HincII, NruI, PvuI and ScaI. The plasmid is called pTrc99AcysDNC (Figure 5).

5b) Preparation of L-threonine with the strain
MG442/pTrc99AcysDNC

- 30 The L-threonine-producing E. coli strain MG442 is described in the patent specification US-A- 4,278,765 and deposited as CMIM B-1628 at the Russian National Collection for Industrial Microorganisms (VKPM, Moscow, Russia).

The strain MG442 is transformed with the expression plasmid pTrc99AcysDNC described in example 5a and with the vector pTrc99A and plasmid-carrying cells are selected on LB agar with 50 µg/ml ampicillin. The strains MG442/pTrc99AcysDNC
5 and MG442/pTrc99A are formed in this manner. Selected individual colonies are then multiplied further on minimal medium with the following composition: 3.5 g/l $\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$, 1.5 g/l KH_2PO_4 , 1 g/l NH_4Cl , 0.1 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 2 g/l glucose, 20 g/l agar, 50 mg/l ampicillin. The formation of
10 L-threonine is checked in batch cultures of 10 ml contained in 100 ml conical flasks. For this, 10 ml of preculture medium of the following composition: 2 g/l yeast extract, 10 g/l $(\text{NH}_4)_2\text{SO}_4$, 1 g/l KH_2PO_4 , 0.5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 15 g/l CaCO_3 , 20 g/l glucose, 50 mg/l ampicillin are inoculated
15 and the batch is incubated for 16 hours at 37°C and 180 rpm on an ESR incubator from Kühner AG (Birsfelden, Switzerland).

250 µl portions of this preculture are transinoculated into 10 ml of production medium (25 g/l $(\text{NH}_4)_2\text{SO}_4$, 2 g/l KH_2PO_4 ,
20 1 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.03 g/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.018 g/l $\text{MnSO}_4 \cdot 1\text{H}_2\text{O}$, 30 g/l CaCO_3 , 20 g/l glucose, 50 mg/l ampicillin) and the batch is incubated for 48 hours at 37°C. The formation of L-threonine by the starting strain MG442 is investigated in the same manner, but no addition of ampicillin to the
25 medium takes place. After the incubation the optical density (OD) of the culture suspension is determined with an LP2W photometer from Dr. Lange (Düsseldorf, Germany) at a measurement wavelength of 660 nm.

The concentration of L-threonine formed is then determined
30 in the sterile-filtered culture supernatant with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column reaction with ninhydrin detection.

The result of the experiment is shown in Table 5.

Table 5

Strain	OD (660 nm)	L-Threonine g/l
MG442	5.6	1.4
MG442/pTrc99A	3.8	1.3
MG442/pTrc99AcysDNC	5.1	2.5

Example 6

Preparation of L-threonine using the *cysJ* and *cysI* genes

5 6a) Construction of the expression plasmid pTrc99AcysJI

The *cysJ* and *cysI* genes from *E. coli* K12 are amplified using the polymerase chain reaction (PCR) and synthetic oligonucleotides. Starting from the nucleotide sequence of the *cysJ* and *cysI* genes in *E. coli* K12 MG1655 (Accession
 10 Number AE000360, Blattner et al. (Science 277: 1453-1462 (1997))), PCR primers are synthesized (MWG Biotech, Ebersberg, Germany):

cysJI1: 5' - CTGGAACATAACGACGCATGAC - 3' (SEQ ID No. 11)

cysJI2: 5' - GACCGGGCTGATGGTTAATCC - 3' (SEQ ID No. 12)

15 The chromosomal *E. coli* K12 MG1655 DNA employed for the PCR is isolated according to the manufacturer's instructions with "Qiagen Genomic-tips 100/G" (QIAGEN, Hilden, Germany). A DNA fragment approx. 3550 bp in size can be amplified with the specific primers under standard PCR conditions
 20 (Innis et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press) with Pfu-DNA polymerase (Promega Corporation, Madison, USA). The PCR product is ligated according to the manufacturer's instructions with

the vector pCR-Blunt II-TOPO (Zero Blunt TOPO PCR Cloning Kit, Invitrogen, Groningen, The Netherlands) and transformed into the E. coli strain TOP10.

Selection of plasmid-carrying cells takes place on LB agar, to which 50 µg/ml kanamycin are added. After isolation of the plasmid DNA, the vector pCR-Blunt II-TOPO-cysJI is cleaved with the restriction enzymes HindIII and XbaI and, after separation in 0.8% agarose gel, the cysJI fragment is isolated with the aid of the QIAquick Gel Extraction Kit (QIAGEN, Hilden, Germany). The vector pTrc99A (Pharmacia Biotech, Uppsala, Sweden) is cleaved with the enzymes HindIII and XbaI and ligation is carried out with the cysJI fragment isolated. The E. coli strain XL1-Blue MRF' (Stratagene, La Jolla, USA) is transformed with the ligation batch and plasmid-carrying cells are selected on LB agar, to which 50 µg/ml ampicillin are added. Successful cloning can be demonstrated after plasmid DNA isolation by control cleavage with the enzymes AccI, ClaI and SphI. The plasmid is called pTrc99AcysJI (Figure 6).

20 6b) Preparation of L-threonine with the strain
MG442/pTrc99AcysJI

The L-threonine-producing E. coli strain MG442 is described in the patent specification US-A- 4,278,765 and deposited as CMIM B-1628 at the Russian National Collection for Industrial Microorganisms (VKPM, Moscow, Russia).

The strain MG442 is transformed with the expression plasmid pTrc99AcysJI described in example 6a and with the vector pTrc99A and plasmid-carrying cells are selected on LB agar with 50 µg/ml ampicillin. The strains MG442/pTrc99AcysJI and MG442/pTrc99A are formed in this manner. Selected individual colonies are then multiplied further on minimal medium with the following composition: 3.5 g/l Na₂HPO₄*2H₂O, 1.5 g/l KH₂PO₄, 1 g/l NH₄Cl, 0.1 g/l MgSO₄*7H₂O, 2 g/l glucose, 20 g/l agar, 50 mg/l ampicillin. The formation of

L-threonine is checked in batch cultures of 10 ml contained in 100 ml conical flasks. For this, 10 ml of preculture medium of the following composition: 2 g/l yeast extract, 10 g/l $(\text{NH}_4)_2\text{SO}_4$, 1 g/l KH_2PO_4 , 0.5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 15 g/l CaCO_3 , 20 g/l glucose, 50 mg/l ampicillin are inoculated and the batch is incubated for 16 hours at 37°C and 180 rpm on an ESR incubator from Kühner AG (Birsfelden, Switzerland).

250 µl portions of this preculture are transinoculated into 10 ml of production medium (25 g/l $(\text{NH}_4)_2\text{SO}_4$, 2 g/l KH_2PO_4 , 1 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.03 g/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.018 g/l $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 30 g/l CaCO_3 , 20 g/l glucose, 50 mg/l ampicillin) and the batch is incubated for 48 hours at 37°C. The formation of L-threonine by the starting strain MG442 is investigated in the same manner, but no addition of ampicillin to the medium takes place. After the incubation the optical density (OD) of the culture suspension is determined with an LP2W photometer from Dr. Lange (Düsseldorf, Germany) at a measurement wavelength of 660 nm.

The concentration of L-threonine formed is then determined in the sterile-filtered culture supernatant with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column reaction with ninhydrin detection.

The result of the experiment is shown in Table 6.

Table 6

Strain	OD (660 nm)	L-Threonine g/l
MG442	5.6	1.4
MG442/pTrc99A	3.8	1.3
MG442/pTrc99AcysJI	6.3	2.5

Example 7

Preparation of L-threonine using the cysH gene

5 7a) Construction of the expression plasmid pTrc99AcysH

The cysH gene from E. coli K12 is amplified using the polymerase chain reaction (PCR) and synthetic oligonucleotides. Starting from the nucleotide sequence of the cysH gene in E. coli K12 MG1655 (Accession Number
10 AE000360, Blattner et al. (Science 277: 1453-1462 (1997))), PCR primers are synthesized (MWG Biotech, Ebersberg, Germany):

cysH1: 5' - GGCAAACAGTGAGGAATCTATG - 3' (SEQ ID No. 13)

cysH2: 5' - GTCCGGCAATATTTACCCTTC - 3' (SEQ ID No. 14)

15 The chromosomal E. coli K12 MG1655 DNA employed for the PCR is isolated according to the manufacturer's instructions with "Qiagen Genomic-tips 100/G" (QIAGEN, Hilden, Germany). A DNA fragment approx. 800 bp in size can be amplified with the specific primers under standard PCR conditions (Innis
20 et al. (1990) PCR Protocols. A Guide to Methods and Applications, Academic Press) with Pfu-DNA polymerase (Promega Corporation, Madison, USA). The PCR product is ligated according to the manufacturer's instructions with

the vector pCR-Blunt II-TOPO (Zero Blunt TOPO PCR Cloning Kit, Invitrogen, Groningen, The Netherlands) and transformed into the E. coli strain TOP10.

Selection of plasmid-carrying cells takes place on LB agar, to which 50 µg/ml kanamycin are added. After isolation of the plasmid DNA, the vector pCR-Blunt II-TOPO-cysH is cleaved with the restriction enzymes HindIII and XbaI and, after separation in 0.8% agarose gel, the cysH fragment is isolated with the aid of the QIAquick Gel Extraction Kit (QIAGEN, Hilden, Germany). The vector pTrc99A (Pharmacia Biotech, Uppsala, Sweden) is cleaved with the enzymes HindIII and XbaI and ligation is carried out with the cysH fragment isolated. The E. coli strain XL1-Blue MRF' (Stratagene, La Jolla, USA) is transformed with the ligation batch and plasmid-carrying cells are selected on LB agar, to which 50 µg/ml ampicillin are added. Successful cloning can be demonstrated after plasmid DNA isolation by control cleavage with the enzymes HincII and MluI. The plasmid is called pTrc99AcysH (Figure 7).

7b) Preparation of L-threonine with the strain MG442/pTrc99AcysH

The L-threonine-producing E. coli strain MG442 is described in the patent specification US-A- 4,278,765 and deposited as CMIM B-1628 at the Russian National Collection for Industrial Microorganisms (VKPM, Moscow, Russia).

The strain MG442 is transformed with the expression plasmid pTrc99AcysH described in example 7a and with the vector pTrc99A and plasmid-carrying cells are selected on LB agar with 50 µg/ml ampicillin. The strains MG442/pTrc99AcysH and MG442/pTrc99A are formed in this manner. Selected individual colonies are then multiplied further on minimal medium with the following composition: 3.5 g/l Na₂HPO₄*2H₂O, 1.5 g/l KH₂PO₄, 1 g/l NH₄Cl, 0.1 g/l MgSO₄*7H₂O, 2 g/l glucose, 20 g/l agar, 50 mg/l ampicillin. The formation of

L-threonine is checked in batch cultures of 10 ml contained in 100 ml conical flasks. For this, 10 ml of preculture medium of the following composition: 2 g/l yeast extract, 10 g/l $(\text{NH}_4)_2\text{SO}_4$, 1 g/l KH_2PO_4 , 0.5 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 15 g/l CaCO_3 , 20 g/l glucose, 50 mg/l ampicillin are inoculated and the batch is incubated for 16 hours at 37°C and 180 rpm on an ESR incubator from Kühner AG (Birsfelden, Switzerland).

250 µl portions of this preculture are transinoculated into 10 ml of production medium (25 g/l $(\text{NH}_4)_2\text{SO}_4$, 2 g/l KH_2PO_4 , 1 g/l $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.03 g/l $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.018 g/l $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 30 g/l CaCO_3 , 20 g/l glucose, 50 mg/l ampicillin) and the batch is incubated for 48 hours at 37°C. The formation of L-threonine by the starting strain MG442 is investigated in the same manner, but no addition of ampicillin to the medium takes place. After the incubation the optical density (OD) of the culture suspension is determined with an LP2W photometer from Dr. Lange (Düsseldorf, Germany) at a measurement wavelength of 660 nm.

The concentration of L-threonine formed is then determined in the sterile-filtered culture supernatant with an amino acid analyzer from Eppendorf-BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column reaction with ninhydrin detection.

The result of the experiment is shown in Table 7.

Table 7

Strain	OD (660 nm)	L-Threonine g/l
MG442	5.6	1.4
MG442/pTrc99A	3.8	1.3
MG442/pTrc99AcysH	4.1	2.7

Brief Description of the Figures:

Figure 1: Map of the plasmid pTrc99AcysB containing the
5 cysB gene.

Figure 2: Map of the plasmid pTrc99AcysK containing the
cysK gene.

Figure 3: Map of the plasmid pTrc99AcysM containing the
cysM gene.

10 Figure 4: Map of the plasmid pTrc99AcysPUWA containing
the cysP, cysU, cysW and cysA genes.

Figure 5: Map of the plasmid pTrc99AcysDNC containing the
cysD, cysN and cysC genes.

Figure 6: Map of the plasmid pTrc99AcysJI containing the
15 cysJ and cysI genes.

Figure 7: Map of the plasmid pTrc99AcysH containing the
cysH gene.

The length data are to be understood as approx. data. The
abbreviations and designations used have the following
20 meaning:

- Amp: Ampicillin resistance gene

- lacI: Gene for the repressor protein of the trc promoter
- Ptrc: trc promoter region, IPTG-inducible
- cysB: Coding region of the cysB gene
- 5 • cysK: Coding region of the cysK gene
- cysM: Coding region of the cysM gene
- cysP: Coding region of the cysP gene
- cysU: Coding region of the cysU gene
- cysW: Coding region of the cysW gene
- 10 • cysA: Coding region of the cysA gene
- cysD: Coding region of the cysD gene
- cysN: Coding region of the cysN gene
- cysC: Coding region of the cysC gene
- cysJ: Coding region of the cysJ gene
- 15 • cysI: Coding region of the cysI gene
- cysH: Coding region of the cysH gene
- 5S: 5S rRNA region
- rrnBT: rRNA terminator region

The abbreviations for the restriction enzymes have the
20 following meaning

- AccI: Restriction endonuclease from *Acinetobacter calcoaceticus*
- BamHI: Restriction endonuclease from *Bacillus amyloliquefaciens* H

- BstEII: Restriction endonuclease from *Bacillus stearothermophilus* ATCC 12980
- ClaI: Restriction endonuclease from *Caryophannon latum*
- EcoRI: Restriction endonuclease from *Escherichia coli*
5 RY13
- EcoRV: Restriction endonuclease from *Escherichia coli*
B946
- HincII: Restriction endonuclease from *Haemophilus influenzae* R_c
- 10 • HindIII: Restriction endonuclease from *Haemophilus influenzae*
- MluI: Restriction endonuclease from *Micrococcus luteus*
IFO 12992
- NdeI: Restriction endonuclease from *Neisseria*
15 *dentrificans*
- NruI: Restriction endonuclease from *Norcadia ruba* (ATCC
15906)
- PauI: Restriction endonuclease from *Paracoccus*
alcaliphilus
- 20 • PvuI: Restriction endonuclease from *Proteus vulgaris*
(ATCC 13315)
- PvuII: Restriction endonuclease from *Proteus vulgaris*
(ATCC 13315)
- ScaI: Restriction endonuclease from *Streptomyces*
25 *caespitosus*
- SmaI: Restriction endonuclease from *Serratia marcescens*

- SpeI: Restriction endonuclease from *Sphaerotilus* species ATCC 13923
- SphI: Restriction endonuclease from *Streptomyces phaeochromogenes*
- 5 • SspI: Restriction endonuclease from *Sphaerotilus* species ATCC 13925
- XbaI: Restriction endonuclease from *Xanthomonas campestris*

What is claimed is:

1. A process for the preparation of L-amino acids, in particular L-threonine, which comprises carrying out the following steps:
 - 5 a) fermentation of microorganisms of the Enterobacteriaceae family which produce the desired L-amino acid and in which at least one or more of the genes of the cysteine biosynthesis pathway chosen from the group consisting of cysG, cysB,
10 cysZ, cysK, cysM, cysA, cysW, cysU, cysP, cysD, cysN, cysC, cysJ, cysI, cysH, cysE and sbp, or nucleotide sequences which code for these, is (are) enhanced, in particular over-expressed,
 - 15 b) concentration of the desired L-amino acid in the medium or in the cells of the microorganisms, and
 - c) isolation of the desired L-amino acid, constituents of the fermentation broth and/or the biomass in its entirety or portions (> 0 to 100%) thereof optionally remaining in the product.
- 20 2. A process as claimed in claim 1, wherein microorganisms in which further genes of the biosynthesis pathway of the desired L-amino acid are additionally enhanced are employed.
- 25 3. A process as claimed in claim 1, wherein microorganisms in which the metabolic pathways which reduce the formation of the desired L-amino acid are at least partly eliminated are employed.
- 30 4. A process as claimed in claim 1, wherein the expression of the polynucleotide (s) which code(s) for one or more of the genes of cysteine biosynthesis chosen from the group consisting of cysG, cysB, cysZ, cysK, cysM, cysA,

cysW, cysU, cysP, cysD, cysN, cysC, cysJ, cysI, cysH, cysE and sbp is increased.

5. A process as claimed in claim 1, wherein the regulatory and/or catalytic properties of the polypeptides (proteins) for which the polynucleotides cysG, cysB, cysZ, cysK, cysM, cysA, cysW, cysU, cysP, cysD, cysN, cysC, cysJ, cysI, cysH, cysE and sbp code are improved or increased.
6. A process as claimed in claim 1, wherein, for the preparation of L-amino acids, microorganisms of the Enterobacteriaceae family in which in addition at the same time one or more of the genes chosen from the group consisting of:
- 6.1 the thrABC operon which codes for aspartate kinase, homoserine dehydrogenase, homoserine kinase and threonine synthase,
 - 6.2 the pyc gene which codes for pyruvate carboxylase,
 - 6.3 the pps gene which codes for phosphoenol pyruvate synthase,
 - 6.4 the ppc gene which codes for phosphoenol pyruvate carboxylase,
 - 6.5 the pntA and pntB genes which code for transhydrogenase,
 - 6.6 the rhtB gene which imparts homoserine resistance,
 - 6.7 the mgo gene which codes for malate:quinone oxidoreductase,
 - 6.8 the rhtC gene which imparts threonine resistance,

- 6.9 the thrE gene which codes for the threonine export protein
- 6.10 the gdhA gene which codes for glutamate dehydrogenase
- 5 6.11 the hns gene which codes for the DNA-binding protein HLP-II,
- 6.12 the pgm gene which codes for phosphoglucomutase,
- 10 6.13 the fba gene which codes for fructose biphosphate aldolase,
- 6.14 the ptsH gene which codes for the phosphohistidine protein hexose phosphotransferase,
- 15 6.15 the ptsI gene which codes for enzyme I of the phosphotransferase system,
- 6.16 the crr gene which codes for the glucose-specific IIA component,
- 6.17 the ptsG gene which codes for the glucose-specific IIBC component,
- 20 6.18 the lrp gene which codes for the regulator of the leucine regulon,
- 6.19 the mopB gene which codes for 10 Kd chaperone,
- 6.20 the ahpC gene which codes for the small sub-unit of alkyl hydroperoxide reductase,
- 25 6.21 the ahpF gene which codes for the large sub-unit of alkyl hydroperoxide reductase,

is or are enhanced, in particular over-expressed, are fermented.

7. A process as claimed in claim 1, wherein, for the preparation of L-amino acids, microorganisms of the Enterobacteriaceae family in which in addition at the same time one or more of the genes chosen from the group consisting of:
- 7.1 the tdh gene which codes for threonine dehydrogenase,
 - 7.2 the mdh gene which codes for malate dehydrogenase,
 - 10 7.3 the gene product of the open reading frame (orf) yjfa,
 - 7.4 the gene product of the open reading frame (orf) yjfp,
 - 15 7.5 the pckA gene which codes for phosphoenolpyruvate carboxykinase,
 - 7.6 the poxB gene which codes for pyruvate oxidase,
 - 7.7 the aceA gene which codes for isocitrate lyase,
 - 7.8 the dgsA gene which codes for the DgsA regulator of the phosphotransferase system,
 - 20 7.9 the fruR gene which codes for the fructose repressor,
 - 7.10 the rpoS gene which codes for the sigma³⁸ factor
- is or are attenuated, in particular eliminated or reduced in expression, are fermented.

Figure 1:




Figure 2:




Figure 3:




Figure 4:




Figure 5:




Figure 6:





Figure 7:



SEQUENCE PROTOCOL

5 <110> Degussa AG
<120> Process for the preparation of L-amino acids using strains of the Enterobacteriaceae family
<130> 010275 BT
10 <160> 14
<170> PatentIn version 3.1
15 <210> 1
<211> 22
<212> DNA
<213> artificial sequence
20 <220>
<221> Primer
<222> (1)..(22)
<223> cysB1
25 <400> 1
gcgtctaagt ggatggttta ac 22
<210> 2
<211> 20
30 <212> DNA
<213> artificial sequence
<220>
<221> Primer
35 <222> (1)..(20)
<223> cysB2
<400> 2
ggtgccgaaa ataacgcaag 20
40 <210> 3
<211> 21
<212> DNA
<213> artificial sequence
45 <220>
<221> Primer
<222> (1)..(21)
<223> cysK1
50 <400> 3
cagttaagga caggccatga g 21
<210> 4
55 <211> 22
<212> DNA
<213> artificial sequence
<220>
60 <221> Primer
<222> (1)..(22)
<223> cysK2
<400> 4
65 gctggcatta ctgttgcaat tc 22

<210> 5
 <211> 28
 <212> DNA
 5 <213> artificial sequence

 <220>
 <221> Primer
 <222> (1)..(28)
 10 <223> cysM1

 <400> 5
 cgcatcagtc tagaccacgt taggatag 28

 15 <210> 6
 <211> 25
 <212> DNA
 <213> artificial sequence

 20 <220>
 <221> Primer
 <222> (1)..(25)
 <223> cysM2

 25 <400> 6
 catcagtcgc cgaagctttt aatcc 25

 <210> 7
 <211> 30
 30 <212> DNA
 <213> artificial sequence

 <220>
 <221> Primer
 35 <222> (1)..(30)
 <223> cysPUWA1

 <400> 7
 gtctctagat aaataagggt gcgcaatggc 30
 40
 <210> 8
 <211> 25
 <212> DNA
 <213> artificial sequence
 45
 <220>
 <221> Primer
 <222> (1)..(25)
 <223> cysPUWA2
 50
 <400> 8
 ccgggcgttt aagcttcact caacc 25

 <210> 9
 55 <211> 30
 <212> DNA
 <213> artificial sequence

 <220>
 <221> Primer
 60 <222> (1)..(30)
 <223> cysDNC1

 <400> 9
 65 gcaagaaaat agcggcttag ataaggaacg 30

<210> 10
 <211> 25
 <212> DNA
 5 <213> artificial sequence

<220>
 <221> Primer
 <222> (1)..(25)
 10 <223> cysDNC2

<400> 10
 catggaaagc ttgtggtgtc tcagg 25

15 <210> 11
 <211> 22
 <212> DNA
 <213> artificial sequence

20 <220>
 <221> Primer
 <222> (1)..(22)
 <223> cysJI1

25 <400> 11
 ctggaacata acgacgcatg ac 22

<210> 12
 <211> 21
 30 <212> DNA
 <213> artificial sequence

<220>
 <221> cysJI2
 35 <222> (1)..(21)
 <223> cysJI2

<400> 12
 gaccgggctg atgggttaatc c 21

40 <210> 13
 <211> 22
 <212> DNA
 <213> artificial sequence

45 <220>
 <221> Primer
 <222> (1)..(22)
 <223> cysH1

50 <400> 13
 ggcaaacagt gaggaatcta tg 22

<210> 14
 55 <211> 21
 <212> DNA
 <213> artificial sequence

<220>
 <221> Primer
 <222> (1)..(21)
 <223> cysH2

<400> 14
 65 gtccggcaat atttaccctt c 21